IL-1beta Murine E. coli

Product Data Sheet

Type: Recombinant
Source: E. coli
Species: Murine
Other names: Catabolin, Lymphocyte-activating factor (LAF), Endogenous Pyrogen (EP), Leukocyte Endogenous Mediator (LEM), Mononuclear Cell Factor (MCF), Interleukin-1beta

Cat. No.:

RBG20001B002	(2 µg)
RBG20001B010	(10 µg)
RBG20001B100	(100 µg)

Description

IL-1beta is a proinflammatory cytokine produced in a variety of cells, including monocytes, tissue macrophages, keratinocytes, and other epithelial cells. Both IL-1alpha and IL-1beta bind to the same receptor and have similar, if not identical, biological properties. These cytokines have a broad range of activities including the stimulation of thymocyte proliferation by inducing IL-2 release, B-cell maturation and proliferation, mitogenic FGF-like activity, and the release of prostaglandin and collagenase from synovial cells. However, whereas IL-1beta is a secreted cytokine, IL-1alpha is predominantly a cell-associated cytokine. Recombinant Murine IL-1beta is a 17.5 kDa protein containing 153 amino acid residues.

Research topic

Cardiovascular disease, Coronary artery disease, Cytokines and chemokines and related molecules, Diabetology - Other Relevant Products, Immune Response, Infection and Inflammation, Neural tissue markers, Oncology, Transplantation

Amino Acid Sequence

```
MVPIRQLHYR LRDEQQKSLV LSDPYELKAL HLNGQNINQQ VIFSMSFVQG EPSNDKIPVA LGLKGKNLYL SCVMKDGTPT
LQLESVDPKQ YPKKKMEKRF VFNKIEVKSK VEFESAEFPN WYISTSQAEH KPVFLGNNSG QDIIDFTMES VSS
```

Source

E. coli

Purity

98%

Biological Activity

The ED_{50} as determined by the dose-dependent stimulation of murine D10S cells is 0.002 ng/ml, corresponding to a specific activity of 5×10^{8} units/mg.

Endotoxin

Endotoxin level is <0.1 ng/µg of protein (<1EU/µg).

Reconstitution

Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

Storage, Stability/Shelf Life

-20°C

HEADQUARTERS: BioVendor Laboratorní medicína, a.s.	Karasek 1767/1	621 00 Brno CZECH REPUBLIC	Phone: +420-549-124-185 Fax: +420-549-211-460	E-mail: Web:	info@biovendor.com sales@biovendor.com www.biovendor.com
AUSTRIA: BioVendor GesmbH	Nußdorfer Straße 20/10	1090 Vienna AUSTRIA	Phone: +43-1-89090-25 Fax: +43-1-89090-2515	E-mail:	infoAustria@biovendor.com
GERMANY, SWITZERLAND: BioVendor GmbH	Otto-Hahn-Straße 16	34123 Kassel GERMANY	Phone: +49-6221-433-9100 Fax: +49-6221-433-9111	E-mail:	infoEU@biovendor.com
USA, CANADA AND MEXICO: BioVendor LLC	128 Bingham Rd. Suite 1300	Asheville, NC 28806 USA	Phone: +1-828-575-9250 +1-800-404-7807 Fax: +1-828-575-9251	E-mail:	infoUSA@biovendor.com