Leucine-Rich Alpha-2-Glycoprotein NATIVE, Canine Serum

Product Data Sheet

Type: Native
Source: Canine pooled serum
Species: Canine
Other names: LRG

Cat. No.:
RD462183050 (0.05 mg)

Description

Native protein isolated from pooled canine serum, 347 AA. MW: 38,129 kDa (calculated without glycosylation). Protein identity confirmed by LC-MS/MS (NCBI no. gi|73987375).

Introduction to the Molecule

Leucine-rich alpha-2-glycoprotein (LRG) is a secretory type 1 acute phase protein whose expression is upregulated by the mediators of acute-phase response. The level of LRG in the blood increases during microbial infections and cancer, presumably as a result of the liver's response to pro-inflammatory cytokines . The function of LRG is unknown but it is proved that LRG is tightly binding to Cyt C. It is suggested that extracellular Cyt C that is released from the cell death by an acute inflammation could be trapped to form tight complex with LRG and be excluded from circulation. LRG secretion by human hepatoma cells is increased by treatment with a mixture of cytokines including IL-6, IL-1b, and TNFalpha that are known to induce acute-phase proteins. It is known that LRG plays a role in adhesive interactions between lymphocytes and the endothelium. LRG may aid in the control of granulopoiesis by helping to modulate the surface expression of different receptor types, including transforming growth factor receptor, granulocyte-macrophage colony-stimulating factor receptor and possibly the macrophage colony stimulating factor receptor. LRG normal plasma concentration is approximately of 21 µg/ml based on purification, or ~50 µg/ml measured by ELISA. Human LRG is a serum glycoprotein of 312 amino acids in length with a predicted unmodified molecular weight of 34 to 36 kDa. SDS PAGE results show LRG molecular weight ranges from 44 to 55 kDa with isoelectric points ranging from 4.52 to 4.72, suggesting that 3 of 30 modifications occur. Leucine-rich alpha-2-glycoprotein consists of a single polypeptide chain with one galactosamine and four glucosamine oligosaccharides attached. The polypeptide has two intrachain disulfide bonds and contains 312 amino acid residues of which 66 are leucine. The amino acid sequence can be exactly divided into 13 segments of 24 residues each, eight of which exhibit a periodic pattern in the occurrence of leucine, proline, and asparagine.

Research topic

Sepsis

Amino Acid Sequence

```
MSPWSPERSQ SPGGLSSHLS KTLLLLLFSV ASAQGVTPDR DTCLVFPSSN GSSVSCHPPA KLPHHFPSDT VHLVVEFFNL
TQLRPDTLQG VSNLQELHLS TNQLESLSPK LLLPVPLLKV LDLTRNALTQ LPPGLFRVSA ALHTLVLKEN RLEVLEPSWL
CGLKALGHLD LSGNHLQTLP PGLLANVTAL RILDLSNNQL KTLPLDLLRG PLQLERLHLE GNRLQVLEEG LLAPQPGLRY
LFLSDNKLAA VAAGAFRGLQ QLDMLDLSNN SLTSVPKGLW TSLGQPTRDM EDGFDISGNP WICDHNLDDL YRWLVANKDK
MFSRNDTRCA RPEALKGQTL LEAAESH
```

Source

Canine pooled serum

Purity

>90%

SDS-PAGE gel



SDS-PAGE analysis of Canine Leucine-rich alpha-2-glycoprotein-1 native protein, 12% gel stained with Coomassie Brillant Blue R250

1) M.W. marker - 14, 21, 31, 45, 66, 97 kDa
2) reduced and boiled sample, 2.5µg/lane
3) non-reduced and non-boiled sample, 2.5µg/lane

Biological Activity

Protein specifically bind to human Cytochrome c1, heme protein - tested.

Endotoxin

< 1.0 EU/ug

Formulation

Filtered (0,4 µm) and lyophilized in 0.5 mg/ml in 20mM TRIS, 50mM NaCl, pH 8.0

Reconstitution

Add deionized water to prepare a working stock solution of approximately 0.5 mg/mL and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

Shipping

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Storage, Stability/Shelf Life

Store lyophilized protein at -20°C. Lyophilized protein remains stable until the expiry date when stored at -20°C. Aliquot reconstituted protein to avoid repeated freezing/thawing cycles and store at -80°C for long term storage. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after one week at 4°C.

Quality Control Test

BCA to determine quantity of the protein.
SDS PAGE to determine purity of the protein.
LAL to determine quantity of endotoxin.

Applications

Cell culture and/or animal studies, ELISA, Immunological methods, Western blotting

Note

This product is intended for research use only.

HEADQUARTERS: BioVendor Laboratorní medicína, a.s.	Karasek 1767/1	621 00 Brno CZECH REPUBLIC	Phone: +420-549-124-185 Fax: +420-549-211-460	E-mail: Web:	info@biovendor.com sales@biovendor.com www.biovendor.com
AUSTRIA: BioVendor GesmbH	Nußdorfer Straße 20/10	1090 Vienna AUSTRIA	Phone: +43-1-89090-25 Fax: +43-1-89090-2515	E-mail:	infoAustria@biovendor.com
GERMANY, SWITZERLAND: BioVendor GmbH	Otto-Hahn-Straße 16	34123 Kassel GERMANY	Phone: +49-6221-433-9100 Fax: +49-6221-433-9111	E-mail:	infoEU@biovendor.com
USA, CANADA AND MEXICO: BioVendor LLC	128 Bingham Rd. Suite 1300	Asheville, NC 28806 USA	Phone: +1-828-575-9250 +1-800-404-7807 Fax: +1-828-575-9251	E-mail:	infoUSA@biovendor.com