

## sRANK Ligand Human E. coli

### Product Data Sheet

<b>Type:</b> Recombinant	<b>Cat. No.:</b>	
<b>Source:</b> E. coli	RBG10283002	(2 µg)
<b>Species:</b> Human	RBG10283010	(10 µg)
<b>Other names:</b> soluble Receptor Activator of NFκB Ligand, TNFSF11, TRANCE (TNF-related activation-induced cytokine), OPGL, ODF (Osteoclast differentiation factor)	RBG10283100	(100 µg)

### Description

RANKL and RANK are members of the TNF superfamily of ligands and receptors that play an important role in the regulation of specific immunity and bone turnover. RANK (receptor) was originally identified as a dendritic cell-membrane protein, which, by interacting with RANKL, augments the ability of dendritic cells. These dendritic cells then stimulate naïve T-cell proliferation in a mixed lymphocyte reaction, promote the survival of RANK + T-cells, and regulate T-cell-dependent immune response. RANKL, which is expressed in a variety of cells, including osteoblasts, fibroblasts, activated T-cells and bone marrow stromal cells, is also capable of interacting with a decoy receptor called OPG. Binding of soluble OPG to sRANKL inhibits osteoclastogenesis by interrupting the signaling between stromal cells and osteoclastic progenitor cells, thereby leading to excess accumulation of bone and cartilage. Human RANKL is reactive on murine cells. Recombinant Human sRANKL is a 20.0 kDa polypeptide comprising the TNF-homologous region of RANKL (176 amino acid residues).

### Research topic

Apoptosis, Bone and cartilage metabolism, Immune Response, Infection and Inflammation

### Amino Acid Sequence

MEKAMVDGSW LDLAKRSKLE AQPFAHLTIN ATDIPSGSHK VSLSSWYHDR GWAKISNMTF SNGKLIVNQD GFYYLYANIC  
FRHHETSGDL ATEYLQLMVY VTKTSIKIPS SHTLMKGGST KYWGSNSEPH FYSINVGFF KLRSGEEISI EVSNPSELLDP  
DQDATYFGAF KVRDID

### Source

E. coli

### Purity

98%

### Biological Activity

Determined by its ability to induce NF-?B in RAW264.7 cells in the absence of any cross-linking. The expected ED<sub>50</sub> for this effect is 10.0-25.0 ng/ml.

### Endotoxin

Endotoxin level is <0.1 ng/µg of protein (<1EU/µg).

### Reconstitution

Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Do not vortex. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

### Storage, Stability/Shelf Life

<b>HEADQUARTERS:</b>				
BioVendor Laboratorní medicína, a.s.	Karasek 1767/1	621 00 Brno CZECH REPUBLIC	Phone: +420-549-124-185 Fax: +420-549-211-460	E-mail: info@biovendor.com sales@biovendor.com Web: www.biovendor.com
AUSTRIA: BioVendor GesmbH	Nußdorfer Straße 20/10	1090 Vienna AUSTRIA	Phone: +43-1-89090-25 Fax: +43-1-89090-2515	E-mail: infoAustria@biovendor.com
GERMANY, SWITZERLAND: BioVendor GmbH	Otto-Hahn-Straße 16	34123 Kassel GERMANY	Phone: +49-6221-433-9100 Fax: +49-6221-433-9111	E-mail: infoEU@biovendor.com
USA, CANADA AND MEXICO: BioVendor LLC	128 Bingham Rd. Suite 1300	Asheville, NC 28806 USA	Phone: +1-828-575-9250 +1-800-404-7807 Fax: +1-828-575-9251	E-mail: infoUSA@biovendor.com