

BioVendor – Laboratorní medicína a.s. Karásek 1767/1, 621 00 Brno, Czech Republic +420 549 124 185 info@biovendor.com sales@biovendor.com www.biovendor.com

WESTERN BLOT PROTOCOL

Annexin A1 Human, Rabbit Polyclonal Antibody

Cat. No.: RD181371100



Western Blot staining of a control protein

MW Marker.: MW: 97, 66, 45, 31, 21, 14 kDa, (Bio-Rad, USA),

	Band of 97 kDa is not showen	
Lane 1:	Recombinant protein	100 ng/Lane, Reducing
Lane 2:	Recombinant protein	10 ng/Lane, Reducing
Lane 3:	Recombinant protein	1 ng/Lane, Reducing
Lane 4:	Recombinant protein	1 ng/Lane, Non-reducing
Lane 5:	Recombinant protein	10 ng/Lane, Non-reducing
Lane 6:	Recombinant protein	100 ng/Lane, Non-reducing

Recombinant Annexin A1 Human E. coli (RD172371100) was subjected to SDS PAGE followed by Western Blot with Annexin A1 Human, Rabbit Polyclonal Antibody (RD181371100) at a concentration of 1 µg/ml. Stained with DAB.



BioVendor – Laboratorní medicína a.s. Karásek 1767/1, 621 00 Brno, Czech Republic +420 549 124 185 info@biovendor.com sales@biovendor.com www.biovendor.com

1. ELFO:

Polyacrylamide gel electrophoresis (PAGE) was used according to the method of Laemmli with minor modifications.

Slab gels (6 x 8 cm), 1 mm thick, were prepared in a multiple gel casting modul (Mini PROTEAN® 3 System, Bio-Rad, USA).

Stacking gel:

4% acrylamide was prepared from a stock solution of 40% acrylamide/bis-acrylamide, 37.5:1 (Bio-Rad, USA) and diluted with 0.8 M Tris (pH 6.8); SDS was added to the final concentration of 0.1%.

Separation gel:

12% polyacrylamide prepared from a stock solution of 40% acrylamide/bis-acrylamide, 37.5:1 (Bio-Rad, USA) and diluted with 1.5 M Tris (pH 8.8); SDS was added to the final concentration of 0.1%.

Polymerisation was achieved with 0.1% v/v N'N'N N-tetramethyl ethylenediamine (TEMED) and 0.1% ammonium persulphate.

Sample preparation:

The protein concentration was determined by the BCA method (with Bovine Albumin as a standard).

Nonreducing conditions:

Protein samples were mixed 1:1 with nonreducing sample buffer (0.19 M Tris, 2% SDS, 1% (v/v) glycerol and 0.001% Bromophenol blue)

Reducing conditions:

Protein samples were mixed 1:1 with reducing sample buffer (0.19 M Tris, 2% SDS, 1% (v/v) glycerol, 0.001% (w/v) Bromophenol blue and 5% 2-Mercaptoethanol) and boiled for 6 min.

Gels were run at 100 V for 15 min and than at 200 V for 45 min. Running Buffer: 0.025 M Tris, 0.192 M glycine and SDS 0.1%, pH 8.3.

2. WESTERN BLOT:

SDS-PAGE separated proteins were blotted onto the PVDF membrane at 15 V for 15 minutes at RT.

<u>Transfer buffer for semidry blotting:</u> 20% methanol, 0.0125 M Tris, 0.096 M glycine and SDS 0.05%.

Membrane with transfered protein was blocked in a blocking buffer for 120 min at RT.

Blocking buffer:

0.05 M Tris, 0.15 M NaCl, 0.1 % Tween, 0.05% Gelatine, 0.02% Thimerosal



BioVendor – Laboratorní medicína a.s. Karásek 1767/1, 621 00 Brno, Czech Republic +420 549 124 185 info@biovendor.com sales@biovendor.com www.biovendor.com

3. DETECTION:

Detection of a Recombinant Annexin A1 Human E. coli (RD172371100) (BioVendor - Laboratorní medicína a.s., Czech Republic)

Primary antibody: **Annexin A1 Human, Rabbit Polyclonal Antibody (RD181371100)** - concentration 1µg/ml in 0.05 M Tris, 0.15 M NaCl, 0.05% Tween, 0.05% Gelatine, 0.02% Thimerosal Incubation: 1 hour Washing: 3x in 0.05 M Tris, 0.15 M NaCl, 0.05% Tween

<u>Secondary antibody:</u> Anti-Rabbit HRP-Conjugate (DAKO) – 1: 2000 in 0.05 M Tris, 0.15 M NaCl, 0.05% Tween, 0.05% Gelatine, 0.02% Thimerosal Incubation: 1 hour Washing: 3x in 0.05 M Tris, 0.15 M NaCl, 0.05% Tween Substrate: DAB