

BioVendor - Laboratorní medicína a.s.

Karásek 1767/1, 621 00 Brno, Czech Republic

+420 549 124 185 info@biovendor.com sales@biovendor.com www.biovendor.com

PRODUCT DATA SHEET

Beta-Microseminoprotein Human, Rabbit Polyclonal Antibody

Cat. No.: RD181140100 Source of Antigen: E. coli Host: Rabbit Size: 0.1 mg Type: Polyclonal Antibody

Other Names:

Prostate secretory protein PSP94, Prostate secreted seminal plasma protein, Prostate secretory protein of 94 amino acids, PSP-94, Seminal plasma beta-inhibin, Immunoglobulin-binding factor, IGBF, PN44, MSMB, PRSP

Preparation:

The antibody was raised in rabbits by immunization with the recombinant Human Beta-microseminoprotein.

Amino Acid Sequence of Immunogen:

The immunization antigen (12.01 kDa – calculated) is a protein containing 104 AA of recombinant Human Beta-microseminoprotein. N-terminal His-tag, 10 extra AA (highlighted).

MKHHHHHAS SCYFIPNEGV PGDSTRKCMD LKGNKHPINS EWQTDNCETC TCYETEISCC TLVSTPVGYD KDNCQRIFKK EDCKYIVVEK KDPKKTCSVS EWII

Purification Method:

Immunoaffinity chromatography on a column with immobilized recombinant Human Beta-microseminoprotein.

Species Reactivity:

Human. Not yet tested in other species.

Antibody Content:

0.1 mg (determined by BCA method, BSA was used as a standard)

Formulation:

The antibody is lyophilized in 0.05 M phosphate buffer, 0.1 M NaCl, pH 7.2. **AZIDE FREE**.

Reconstitution:

Add 0.2 ml of deionized water and let the lyophilized pellet dissolve completely. Slight turbidity may occur after reconstitution, which does not affect activity of the antibody. In this case clarify the solution by centrifugation.

Shipping:

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Storage/Stability:

The lyophilized antibody remains stable and fully active until the expiry date when stored at -20° C.Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles and store frozen at -80° C. Reconstituted antibody can be stored at 4°C for a limited period of time; it does not show decline in activity after one week at 4°C.

Quality Control:

Indirect ELISA – to determine titer of the antibody SDS PAGE – to determine purity of the antibody BCA - to determine quantity of the antibody

Applications:

ELISA, Immunohistochemistry, Western blotting

Note:

This product is for research use only.