

PRODUCT DATA SHEET

Angiopoietin-Like Protein 4 Human, Rabbit Polyclonal Antibody

Cat. No.: RD181073100-01**Size:** 0.1 mg**Source of Antigen:** E. coli**Type:** Polyclonal Antibody**Host:** Rabbit**Other Names:**

Angiopoietin-related protein 4, Hepatic fibrinogen/angiopoietin-related protein, HFARP, ANGPTL4, ARP4, HFARP, PGAR, PP1158, PSEC0166, UNQ171/PRO197, FIAF

Preparation:

The antibody was raised in rabbits by immunization with the recombinant Human ANGPTL4.

Amino Acid Sequence of Immunogen:

The immunization antigen (25 kDa) is a protein containing 220 AA of recombinant Human ANGPTL4. N-Terminal His-tag, 16 extra AA.

MRGSHHHHHH GMASHMGPVQ SKSPRFASWD EMNVLAHGLL QLGQGLREHA ETRTSQLSAL
ERRLSACGSA CQGTEGSTDL PLAPESRVDP EVLHSLQTQL KAQNSRIQQL FHKVAQQQRH
LEKQHLRIQH LQSQFGLLDH KHLDEHAVKP ARRKRLPEMA QPVDPAHNVS RLHRLPRDCQ
ELFQVGERQS GLFEIQPQGS PPFLVNCKMT SDGGWTVIQR

The amino acid sequence of the recombinant Human ANGPTL4 is 100% homologous to the amino acid sequence of the Human ANGPTL4 precursor (AA 26–229) without signal sequence

Purification Method:

Immunoaffinity chromatography on a column with immobilized recombinant Human ANGPTL4.

Species Reactivity:

Human. Not yet tested in other species.

Antibody Content:

0.1 mg (determined by BCA method, BSA was used as a standard)

Formulation:

The antibody is lyophilized in 0.05 M phosphate buffer, 0.1 M NaCl, pH 7.2.

Reconstitution:

Add 0.2 ml of deionized water and let the lyophilized pellet dissolve completely. Slight turbidity may occur after reconstitution, which does not affect activity of the antibody. In this case clarify the solution by centrifugation.

Shipping:

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Storage/Stability:

The lyophilized antibody remains stable and fully active until the expiry date when stored at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles and store frozen at -80°C. Reconstituted antibody can be stored at 4°C for a limited period of time; it does not show decline in activity after one week at 4°C.

Quality Control:

SDS PAGE - to determine purity of the antibody

BCA - to determine quantity of the antibody

Applications:

ELISA, Immunohistochemistry, Western blotting

Note:

This product is for research use only.