

BioVendor - Laboratorní medicína a.s.

Karásek 1767/1, 621 00 Brno, Czech Republic +420 549 124 185 info@biovendor.com sales@biovendor.com www.biovendor.com

PRODUCT DATA SHEET

Leucine-Rich Alpha-2-Glycoprotein Human, Sheep Polyclonal Antibody

Cat. No.: RD184183100 **Size:** 0.1 mg

Source of Antigen: Human pooled serum Type: Polyclonal Antibody

Host: Sheep

Other Names:

LRG

Preparation:

The antibody was raised in sheep by immunization with the native Human LRG.

Amino Acid Sequence of Immunogen:

Native protein isolated from pooled human serum, 312 AA. MW: 34.35 kDa (calculated without glycosylation). Protein identity confirmed by LC-MS/MS (NCBI no. gi|16418467).

VTLSPKDCQV FRSDHGSSIS CQPPAEIPGY LPADTVHLAV EFFNLTHLPA NLLQGASKLQ ELHLSSNGLE SLSPEFLRPV PQLRVLDLTR NALTGLPPGL FQASATLDTL VLKENQLEVL EVSWLHGLKA LGHLDLSGNR LRKLPPGLLA NFTLLRTLDL GENQLETLPP DLLRGPLQLE RLHLEGNKLQ VLGKDLLLPQ PDLRYLFLNG NKLARVAAGA FQGLRQLDML DLSNNSLASV PEGLWASLGQ PNWDMRDGFD ISGNPWICDQ NLSDLYRWLQ AQKDKMFSQN DTRCAGPEAV KGQTLLAVAK SQ

Purification Method:

Immunoaffinity chromatography on a column with immobilized native Human LRG.

Species Reactivity:

Human. Not yet tested in other species.

Antibody Content:

0.1 mg (determined by BCA method, BSA was used as a standard)

Formulation:

The antibody is lyophilized in 0.05 M phosphate buffer, 0.1 M NaCl, pH 7.2. **AZIDE FREE**.

Reconstitution:

Add 0.2 ml of deionized water and let the lyophilized pellet dissolve completely. Slight turbidity may occur after reconstitution, which does not affect activity of the antibody. In this case clarify the solution by centrifugation.

Shipping:

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Date of Issue: 19.05.2023 Page:1 / 2

Storage/Stability:

The lyophilized antibody remains stable and fully active until the expiry date when stored at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles and store frozen at -80°C. Reconstituted antibody can be stored at 4°C for a limited period of time; it does not show decline in activity after one week at 4°C.

Quality Control:

Indirect ELISA – to determine titer of the antibody SDS PAGE – to determine purity of the antibody BCA - to determine quantity of the antibody

Applications:

Immunohistochemistry, Western blotting

Note:

This product is for research use only.

Date of Issue: 19.05.2023 Page:2 / 2