

BioVendor - Laboratorní medicína a.s.

Karásek 1767/1, 621 00 Brno, Czech Republic +420 549 124 185 info@biovendor.com sales@biovendor.com www.biovendor.com

PRODUCT DATA SHEET

Zymogen Granule Membrane Protein 16 Human, Rabbit Polyclonal Antibody

Cat. No.: RD181229100 **Size:** 0.1 mg

Source of Antigen: E. coli Type: Polyclonal Antibody

Host: Rabbit

Other Names:

ZG16, Secretory lectin ZG16

Preparation:

The antibody was raised in rabbits against recombinant Zymogen Granule Membrane Protein 16.

Amino Acid Sequence of Immunogen:

The immunization antigen (17.9 kDa) is a protein containing 161 AA of recombinant Human Zymogen Granule Membrane Protein 16. N-terminal His-tag, 10 extra AA.

MKHHHHHHAS NAIQARSSSY SGEYGSGGGK RFSHSGNQLD GPITALRVRV NTYYIVGLQV RYGKVWSDYV GGRNGDLEEI FLHPGESVIQ VSGKYKWYLK KLVFVTDKGR YLSFGKDSGT SFNAVPLHPN TVLRFISGRS GSLIDAIGLH WDVYPTSCSR C

Purification Method:

Immunoaffinity chromatography on a column with immobilized recombinant Human Zymogen Granule Membrane Protein 16.

Species Reactivity:

Human. Not yet tested in other species.

Antibody Content:

0.1 mg (determined by BCA method, BSA was used as a standard)

Formulation:

The antibody is lyophilized in 0.05 M phosphate buffer, 0.1 M NaCl, pH 7.2. **AZIDE FREE**.

Reconstitution:

Add 0.2 ml of deionized water and let the lyophilized pellet dissolve completely. Slight turbidity may occur after reconstitution, which does not affect activity of the antibody. In this case clarify the solution by centrifugation.

Shipping:

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Date of Issue: 19.05.2023 Page:1 / 2

Storage/Stability:

The lyophilized antibody remains stable and fully active until the expiry date when stored at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles and store frozen at -80°C. Reconstituted antibody can be stored at 4°C for a limited period of time; it does not show decline in activity after one week at 4°C.

Quality Control:

Indirect ELISA – to determine titer of the antibody SDS PAGE – to determine purity of the antibody BCA - to determine quantity of the antibody

Applications:

ELISA, Immunohistochemistry, Western blotting

Note:

This product is for research use only.

Date of Issue: 19.05.2023 Page:2 / 2