

## Interleukin-9 Rat E. coli

### Product Data Sheet

|   |                  |          |
|---|------------------|----------|
| <b>Type:</b> Recombinant  | <b>Cat. No.:</b> |          |
| <b>Source:</b> E. coli  | RBG30009002      | (2 µg)   |
| <b>Species:</b> Rat   | RBG30009010      | (10 µg)  |
| <b>Other names:</b> IL-9, MEA, P40 cytokine, P40 T-cell and mast cell growth factor, T-cell/mast cell growth factor P40, TCGF III | RBG30009100      | (100 µg) |

### Description

IL-9 is an immunoregulatory cytokine produced by IL-2 activated Th2 lymphocytes. IL-9 enhances the proliferation of T lymphocytes, mast cells, erythroid precursor cells and megakaryoblastic leukemia cell lines. Over-expression of IL-9 has been implicated in the pathogenesis of anaplastic lymphoma and Hodgkin's disease. Recombinant Rat IL-9 is a 14.3 kDa protein of 127 amino acid residues, including 10 cysteine residues that are fully conserved between the human and murine proteins.

### Introduction to the Molecule

Interleukin-9 (IL-9) is a proinflammatory cytokine historically believed to be involved in type 2 immune responses. However, recent evidence suggests IL-9 may be secreted by other T Helper lineages such as Treg and Th17 in addition to a new category called Th9. This Th9 lineage can either be derived from Th2 cells with TGF beta or differentiated directly from naïve CD4+ T cells with TGF beta and IL-4. The IL-9 expression in subsets such as Treg and Th17 illustrates the plasticity of cells to reprogram to alternative fates. IL-9 is a member of the common cytokine receptor gamma chain-dependent family of cytokines which also includes IL-2, IL-4, IL-7, IL-15 and IL-21. IL-9 is an extensively glycosylated protein of 14 kDa containing ten cysteine residues involved in disulfide bonding. The human gene maps to chromosome 5q31-32 which is a chromosomal region sometimes deleted in patients with myelodysplastic syndrome. Its pleiotropic effects on Th2 lymphocytes, B lymphocytes, mast cells, eosinophils, IgE production and gut and airway epithelial cells have implicated IL-9 in asthma and other allergy-related disorders. The existence of an IL-9- mediated autocrine loop has been suggested for some malignancies such as Hodgkin's disease and large cell anaplastic lymphoma for Hodgkin's cell lines. IL-9 is expressed by Reed-Sternberg cells and Hodgkin lymphoma cells and some large aplastic lymphoma cells, while non-Hodgkin lymphomas and peripheral T-cell lymphomas do not express it

### Research topic

Cytokines and chemokines and related molecules

### Amino Acid Sequence

MQRCTSWG I QHTSYLIENL KDDPSSKCSC SANVTSLCL PIPSDDCTTP CFQEGMSQVT NATQQSKFSP FFFRVKRIVE  
TLKSNKCQFF SCEKPCNQTT AGNTVSFLKS LLKTFQKTEV QVQRSRA

### Source

E. coli

### Purity

98%

### Biological Activity

Data not available.

### Endotoxin

Endotoxin level is <0.1 ng/µg of protein (<1EU/µg).

### Reconstitution

Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Do not vortex. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

## Storage, Stability/Shelf Life

-20°C

|  |                               |                               |   |   |
|--|-------------------------------|-------------------------------|---|---|
| <b>HEADQUARTERS:</b>                     |                               |                               |   |   |
| BioVendor Laboratorní<br>medicína, a.s.  | Karasek 1767/1                | 621 00 Brno<br>CZECH REPUBLIC | Phone: +420-549-124-185<br>Fax: +420-549-211-460                  | E-mail: info@biovendor.com<br>sales@biovendor.com<br>Web: www.biovendor.com |
| AUSTRIA:<br>BioVendor GesmbH             | Nußdorfer Straße 20/10        | 1090 Vienna<br>AUSTRIA        | Phone: +43-1-89090-25<br>Fax: +43-1-89090-2515                    | E-mail: infoAustria@biovendor.com   |
| GERMANY, SWITZERLAND:<br>BioVendor GmbH  | Otto-Hahn-Straße 16           | 34123 Kassel<br>GERMANY       | Phone: +49-6221-433-9100<br>Fax: +49-6221-433-9111                | E-mail: infoEU@biovendor.com  |
| USA, CANADA AND MEXICO:<br>BioVendor LLC | 128 Bingham Rd.<br>Suite 1300 | Asheville, NC 28806<br>USA    | Phone: +1-828-575-9250<br>+1-800-404-7807<br>Fax: +1-828-575-9251 | E-mail: infoUSA@biovendor.com   |