

Interleukin-4 Murine E. coli

Product Data Sheet

Type: Recombinant	Cat. No.:	
Source: E. coli	RBG20004005	(5 µg)
Species: Murine	RBG20004020	(20 µg)
Other names: IL-4, B-cell stimulatory factor 1, Lymphocyte stimulatory factor 1, Binetrakin, Pitrakinra, B cell differentiation factor gamma, B cell growth factor	RBG20004100	(100 µg)

Description

IL-4 is a pleiotropic cytokine that regulates diverse T and B cell responses including cell proliferation, survival and gene expression. Produced by mast cells, T cells and bone marrow stromal cells, IL-4 regulates the differentiation of naive CD4+ T cells into helper Th2 cells, characterized by their cytokine-secretion profile that includes secretion of IL-4, IL-5, IL-6, IL-10, and IL-13, which favor a humoral immune response. Another dominant function of IL-4 is the regulation of immunoglobulin class switching to the IgG1 and IgE isotypes. Excessive IL-4 production by Th2 cells has been associated with elevated IgE production and allergy. Recombinant Murine IL-4 is a 13.5 kDa globular protein containing 121 amino acid residues.

Introduction to the Molecule

IL-4 (15 to 19 kDa) is produced by a variety of T cell thymocyte populations, as well as some mast cell precursors that are non-T or B cell types. IL-4 plays an important role in the activation of resting B-cells and in the subsequent proliferation and differentiation of B-cells. IL-4 is essential for IgE synthesis in vitro and its involvement with allergic diseases has been investigated. IL-4 has anti-inflammatory effects.

Research topic

Animal studies, Cytokines and chemokines and related molecules

Amino Acid Sequence

MHIHGCDKNH LREIIGILNE VTGEGTPCTE MDVPNVLAT KNTTESELVC RASKVLRIFY LKHGKTPCLK KNSSVLMELQ
RLFRAFRCILD SSISCTMNES KSTSLKDFLE SLKSIMQMDY S

Source

E. coli

Purity

98%

Biological Activity

The ED₅₀ was determined by the dose-dependent proliferation of murine HT-2 cells is ? 2.0 ng/ml, corresponding to a specific activity of ? 5x10⁵ units/mg.

Endotoxin

Endotoxin level is <0.1 ng/µg of protein (<1EU/µg).

Reconstitution

Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

Storage, Stability/Shelf Life

HEADQUARTERS:				
BioVendor Laboratorní medicína, a.s.	Karasek 1767/1	621 00 Brno CZECH REPUBLIC	Phone: +420-549-124-185 Fax: +420-549-211-460	E-mail: info@biovendor.com sales@biovendor.com Web: www.biovendor.com
AUSTRIA: BioVendor GesmbH	Nußdorfer Straße 20/10	1090 Vienna AUSTRIA	Phone: +43-1-89090-25 Fax: +43-1-89090-2515	E-mail: infoAustria@biovendor.com
GERMANY, SWITZERLAND: BioVendor GmbH	Otto-Hahn-Straße 16	34123 Kassel GERMANY	Phone: +49-6221-433-9100 Fax: +49-6221-433-9111	E-mail: infoEU@biovendor.com
USA, CANADA AND MEXICO: BioVendor LLC	128 Bingham Rd. Suite 1300	Asheville, NC 28806 USA	Phone: +1-828-575-9250 +1-800-404-7807 Fax: +1-828-575-9251	E-mail: infoUSA@biovendor.com