

## Interleukin-22 Human E. coli

### Product Data Sheet

<b>Type:</b> Recombinant	<b>Cat. No.:</b>	
<b>Source:</b> E. coli	RBG10022002	(2 µg)
<b>Species:</b> Human	RBG10022010	(10 µg)
<b>Other names:</b> IL-22, ILD110, ILTIF, Interleukin 10 related T cell derived inducible factor, TIFIL23, TIFa, zcyto18	RBG10022100	(100 µg)

### Description

IL-22 is a member of the IL-10 family of regulatory cytokines, which includes IL-10, IL-19, IL-20, IL-22, IL-24 and IL-26. Members of this family share partial homology in their amino acid sequences, but they are dissimilar in their biological functions. Produced by T lymphocytes, IL-22 inhibits IL-4 production by Th2 cells, and induces acute phase reactants in the liver and pancreas. IL-22 signals through a receptor system consisting of IL-10Rbeta/CRF2-4 and IL-22R, both of which are members of the class II cytokine-receptor family. Recombinant Human IL-22 is a 33.6 kDa, non-disulfide-linked, homodimeric protein consisting of two 147 amino acid polypeptide chains.

### Introduction to the Molecule

Interleukin-22 is a new cytokine that regulates the production of acute phase proteins of the immunological response. It is produced by activated Th1 and NK cells acting primarily on epithelial cells and is involved in inflammatory responses. IL-22 plays a protective role in T cell-mediated hepatitis induced by Concanavalin A, acting as a survival factor for hepatocytes. IL-22 is present in high quantities in the blood of Crohn's disease patients in contrast to IFN-gamma and IL-17 levels.

### Research topic

Animal studies, Cytokines and chemokines and related molecules

### Amino Acid Sequence

MAPISSHCRL DKSNFQQPYI TNRTFMLAKE ASLADNNTDV RLIGEKLFHG VMSERCYLM KQVLNFTLEE VLFPPQSDRFQ  
PYMQEVPVFL ARLSNRLSTC HIEGDDLHIQ RNVQKLKDTV KKLGESGEIK AIGELDLLFM SLRNACI

### Source

E. coli

### Purity

98%

### Biological Activity

Assay #1: Determined by its ability to activate STAT following receptor ligand interaction. Assay #2: Measured by its ability to induce IL-10 secretion in COLO 205 (human colon carcinoma cells). The expected ED<sub>50</sub> for this effect is 0.7- 1.0 ng/ml.

### Endotoxin

Endotoxin level is <0.1 ng/µg of protein (<1EU/µg).

### Reconstitution

Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.5-1.0 mg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

### Storage, Stability/Shelf Life

<b>HEADQUARTERS:</b> BioVendor Laboratorní medicína, a.s.	Karasek 1767/1	621 00 Brno CZECH REPUBLIC	Phone: +420-549-124-185 Fax: +420-549-211-460	E-mail: info@biovendor.com sales@biovendor.com Web: www.biovendor.com
AUSTRIA: BioVendor GesmbH	Nußdorfer Straße 20/10	1090 Vienna AUSTRIA	Phone: +43-1-89090-25 Fax: +43-1-89090-2515	E-mail: infoAustria@biovendor.com
GERMANY, SWITZERLAND: BioVendor GmbH	Otto-Hahn-Straße 16	34123 Kassel GERMANY	Phone: +49-6221-433-9100 Fax: +49-6221-433-9111	E-mail: infoEU@biovendor.com
USA, CANADA AND MEXICO: BioVendor LLC	128 Bingham Rd. Suite 1300	Asheville, NC 28806 USA	Phone: +1-828-575-9250 +1-800-404-7807 Fax: +1-828-575-9251	E-mail: infoUSA@biovendor.com