

Interleukin-13 Murine E. coli

Product Data Sheet

Type: Recombinant	Cat. No.:	
Source: E. coli	RBG20013002	(2 µg)
Species: Murine	RBG20013010	(10 µg)
Other names: IL-13, ALRH	RBG20013100	(100 µg)

Description

IL-13 is an immunoregulatory cytokine produced primarily by activated Th2 cells, and also by mast cells and NK cells. Targeted deletion of IL-13 in mice resulted in impaired Th2 cell development and indicated an important role for IL-13 in the expulsion of gastrointestinal parasites. IL-13 exerts anti-inflammatory effects on monocytes and macrophages and it inhibits the expression of inflammatory cytokines such as IL-1beta, TNF-alpha, IL-6 and IL-8. IL-13 has also been shown to enhance B cell proliferation and to induce isotype switching, resulting in increased production of IgE. Blocking of IL-13 activity inhibits the pathophysiology of asthma. Human and murine IL-13 are cross-species reactive. Recombinant Murine IL-13 is a 12.3 kDa protein consisting of 111 amino acid residues.

Introduction to the Molecule

Interleukin-13 (IL-13) is a Th2-type cytokine, secreted from CD4(+) T cells, mast cells, basophils and eosinophils. It is an immunoregulatory cytokine. It has become evident that IL-13 is a key mediator in the pathogenesis of allergic inflammation. The important role of IL-13 in the pathogenesis of bronchial asthma as well as other allergic diseases has been recognized. Additionally, IL-13 plays a major role in various other inflammatory diseases including cancer.

Research topic

Animal studies, Cytokines and chemokines and related molecules

Amino Acid Sequence

MPVPRSVSLP LTLKELIEEL SNITQDQTPL CNGSMVWSVD LAAGGFCVAL DSLTNISNCN AIYRTQRILH GLCNRKAPTT
VSSLPDTKIE VAHFITKLLS YTKQLFRHGP F

Source

E. coli

Purity

98%

Biological Activity

The ED₅₀ as determined by the dose-dependent proliferation of TF-1 cells was 4.0 ng/ml, corresponding to a specific activity of 2.5x10⁵ units/mg.

Endotoxin

Endotoxin level is <0.1 ng/µg of protein (<1EU/µg).

Reconstitution

Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Note: Slow to dissolve. Do not vortex. This solution can be stored at 2-8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

Storage, Stability/Shelf Life

HEADQUARTERS:				
BioVendor Laboratorní medicína, a.s.	Karasek 1767/1	621 00 Brno CZECH REPUBLIC	Phone: +420-549-124-185 Fax: +420-549-211-460	E-mail: info@biovendor.com sales@biovendor.com Web: www.biovendor.com
AUSTRIA: BioVendor GesmbH	Nußdorfer Straße 20/10	1090 Vienna AUSTRIA	Phone: +43-1-89090-25 Fax: +43-1-89090-2515	E-mail: infoAustria@biovendor.com
GERMANY, SWITZERLAND: BioVendor GmbH	Otto-Hahn-Straße 16	34123 Kassel GERMANY	Phone: +49-6221-433-9100 Fax: +49-6221-433-9111	E-mail: infoEU@biovendor.com
USA, CANADA AND MEXICO: BioVendor LLC	128 Bingham Rd. Suite 1300	Asheville, NC 28806 USA	Phone: +1-828-575-9250 +1-800-404-7807 Fax: +1-828-575-9251	E-mail: infoUSA@biovendor.com