

Thyrostimulin alpha subunit (Glycoprotein hormone alpha 2) Human E. coli

Product Data Sheet

Type: Recombinant

Source: E. coli

Species: Human

Other names: Thyrostimulin alpha subunit, GPHA2, Glycoprotein hormone beta 5, GPHB5, Thyrostimulin beta subunit, Glycoprotein hormone alpha-2, Putative secreted protein Zsig51, GPA2, ZSIG51

Cat. No.:

RD172110100

(0.1 mg)

Description

Total 120 AA. Mw: 13.28 kDa (calculated). UniProtKB acc.no. Q96T91. N-terminal His-tag (14AA).

Introduction to the Molecule

Human thyrostimulin ranks among the glycoprotein hormone family. These hormones consist of two subunits, the common alpha- and specific beta-subunits, which associate noncovalently to form a heterodimer. The alpha-subunit combines with four distinct beta-subunits giving rise to four biologically active hormones in human: FSH, LH, TSH, and CG. FSH, LH, and TSH, mainly expressed in the anterior pituitary, are essential for coordinated endocrine regulation in the hypothalamus-pituitary axis and show to activate specific G protein-coupled receptors in the thyroid (TSH receptor) and gonads (LH and FSH receptors), respectively.

The heterodimeric glycoprotein hormones have only been identified in vertebrates and are highly conserved in organisms from primitive rayfin fish (Chondrostei) to human in both primary sequences and functional characteristics.

Corticotroph-derived glycoprotein hormone (CGH), also referred to as thyrostimulin, is a noncovalent heterodimer of glycoprotein hormone alpha 2 (GPHA2) and glycoprotein hormone beta 5 (GPHB5). Recombinant A2/B5 heterodimeric glycoproteins activates human TSH receptors, but not LH and FSH receptors, and shows high affinity to TSH receptors in a radioligand receptor assay. The heterodimer also stimulates cAMP production and thymidine incorporation by cultured thyroid cells and increases serum thyroxine levels in TSH-suppressed rats in vivo. This new heterodimeric glycoprotein hormone was named as thyrostimulin based on its thyroid-stimulating activity. The expression of thyrostimulin in the anterior pituitary known to express TSH receptors suggested a paracrine mechanism.

Research topic

Others

Amino Acid Sequence

MRGSHHHHHH GMASQEAVIP GCHLHPFNVT VRSDRQGTCQ GSHVAQACVG HCESSAFPSR YSVLVASGYR HNITSVSVQCC
TISGLKKVKV QLQCVGSRRE ELEIFTARAC QCDMCLRSRY


Source

E. coli

Purity

>95%

SDS-PAGE gel



12% SDS-PAGE separation of Human Thyrostimulin alpha subunit

1. M.W. marker - 14, 21, 31, 45, 66, 97 kDa
2. reduced and heated sample, 5µg/lane
3. non-reduced and non-heated sample, 5µg/lane

Formulation

Filtered (0.4 µm) and lyophilized in 0.5 mg/mL in 0.05 M Acetate buffer pH=4.0

Reconstitution

Add 0.1M Acetate buffer pH=4.0 to prepare a working stock solution of approximately 0.5 mg/mL and let the lyophilized pellet dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 10µg/mL. In higher concentrations the solubility of this antigen is limited. Filter sterilize your culture media/working solutions containing this non-sterile product before using in cell culture.

Shipping

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Storage, Stability/Shelf Life

Store lyophilized protein at -80°C. Lyophilized protein remains stable until the expiry date when stored at -80°C. Aliquot reconstituted protein to avoid repeated freezing/thawing cycles and store at -80°C for long term storage. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

Quality Control Test

BCA to determine quantity of the protein.
SDS PAGE to determine purity of the protein.

Applications

Western blotting

Note

This product is intended for research use only.

HEADQUARTERS: BioVendor Laboratorní medicína, a.s.	Karasek 1767/1	621 00 Brno CZECH REPUBLIC	Phone: +420-549-124-185 Fax: +420-549-211-460	E-mail: info@biovendor.com sales@biovendor.com Web: www.biovendor.com
AUSTRIA: BioVendor GesmbH	Nußdorfer Straße 20/10	1090 Vienna AUSTRIA	Phone: +43-1-89090-25 Fax: +43-1-89090-2515	E-mail: infoAustria@biovendor.com
GERMANY, SWITZERLAND: BioVendor GmbH	Otto-Hahn-Straße 16	34123 Kassel GERMANY	Phone: +49-6221-433-9100 Fax: +49-6221-433-9111	E-mail: infoEU@biovendor.com
USA, CANADA AND MEXICO: BioVendor LLC	128 Bingham Rd. Suite 1300	Asheville, NC 28806 USA	Phone: +1-828-575-9250 +1-800-404-7807 Fax: +1-828-575-9251	E-mail: infoUSA@biovendor.com